


Ocrevus Prescription / Enrollment Form	Y-Medical Associates Fax Referral To: 855-838-0623 Phone: 800-447-7558	
Date: _____		

PATIENT INFORMATION	PRESCRIBER INFORMATION
Patient Name: _____ Address: _____ City, State, Zip: _____ Home Phone: _____ Cell Phone: _____ Alternate Phone: _____ Date of Birth: _____ Gender: <input type="checkbox"/> M <input type="checkbox"/> F	Prescriber Name: _____ Address: _____ City, State, Zip: _____ Phone: _____ Fax: _____ DEA#: _____ NPI#: _____ Contact Person: _____

INSURANCE INFORMATION (Please attach the front and back of insurance and prescription drug card)					
Primary Insurance: _____	ID#: _____	Group: _____			
Secondary Insurance: _____	ID#: _____	Group: _____			
Prescription Card: _____	ID#: _____	BIN: _____	PCN: _____	Group: _____	

DIAGNOSIS (ICD-10)	
<input type="checkbox"/> G35 Relapsing forms of Multiple Sclerosis (Clinically isolated syndrome/relapsing-remitting disease/active secondary progressive disease) <input type="checkbox"/> G35 Primary Progressive Multiple Sclerosis	

PRE-SCREENING	
<input type="checkbox"/> Hepatitis B Surface Antigen: _____ <input type="checkbox"/> Total Hepatitis B Core Antibody (Anti-HBc): _____ <input type="checkbox"/> Serum Immunoglobulins: _____ <input type="checkbox"/> Vaccination: _____ <small>(live or live-attenuated 4 weeks before, non-live 2 weeks before initiation of therapy)</small> Labs (During Therapy): _____	Vaccinations: Live-attenuated or live vaccines is not recommended during treatment and after discontinuation until B-cell repletion. Administer all necessary immunizations according to immunization guidelines at least 4 weeks prior to initiation for live or attenuated vaccines and at least 2 weeks prior to initiation for non-live vaccines. Pre-screening: Required Hepatitis screening before first dose to include: ____ Hepatitis B Surface Antigen (HBsAg) and Total Hepatitis B Core Antibody (anti-HBc) * Ocrevus® is contraindicated in patients with active HBV. Patients who are negative for surface antigen HBsAg (-) and positive for HB core antibody HBcAB (+) or positive for surface antigen HBsAg (+), should consult liver disease experts before starting and during treatment. ____ Quantitative Serum Immunoglobulin Screening (IgG, IgA, IgM)

PRESCRIPTION ORDERS	
Premeds Premedication to be given 30 minutes prior to infusion: <input type="checkbox"/> Acetaminophen PO: <input type="checkbox"/> 325mg <input type="checkbox"/> 500mg <input type="checkbox"/> 650mg Diphenhydramine: <input type="checkbox"/> 25mg IVP <input type="checkbox"/> 50mg IVP <input type="checkbox"/> 25mg PO <input type="checkbox"/> 50mg PO OR <input type="checkbox"/> Alternate oral antihistamine: <input type="checkbox"/> Cetirizine 10mg <input type="checkbox"/> Loratadine 10mg <input type="checkbox"/> Fexofenadine 60mgs <input type="checkbox"/> Fexofenadine 180mgs IV Access Flush Order: NaCl 0.9% 5-10ml IV before and after infusion <input type="checkbox"/> Methylprednisolone <input type="checkbox"/> 125mg IVP <input type="checkbox"/> 40mg IVP OR <input type="checkbox"/> _____mg PO <input type="checkbox"/> Others/Miscellaneous: _____	Anaphylaxis Orders and Medications Diphenhydramine Administer 25 mg slow IV/IM may repeat x1 Dispense: 1 x 50 mg vial Epinephrine <input type="checkbox"/> Administer 0.15mg (1:2000) Sub-Q (< 30 Kg) <input type="checkbox"/> Administer 0.3mg (1:1000) Sub-Q (≥ 30 Kg) Dispense: 1 package Sodium Chloride 0.9% <i>Use to maintain IV line, prevent or treat hypotension in case of anaphylaxis</i> Dispense: QS

Medication	
Ocrevus (Ocrelizumab) IV as directed to infuse per protocol via pump with 0.22 µ [1:100] filter, following each infusion with a one hour post observation period. <input type="checkbox"/> Induction/Initial dosing: Induction/Initial dosing: 300mg Ocrevus IV in 250ml Sodium Chloride 0.9% to be infused at Week 0 over 2.5 hours or longer and 2 weeks later over 2.5 hours or longer. No Refills. **To be infused in MD office or an Infusion suite. <input type="checkbox"/> Maintenance dosing: 600mg Ocrevus IV in 500ml Sodium Chloride 0.9% to be infused every 6 months. <input type="checkbox"/> 2 hrs or longer for eligible patients who have not experienced a serious infusion reaction with any previous Ocrevus Infusion <input type="checkbox"/> 3.5-4 hrs or longer. Refills: <input type="checkbox"/> X1 year **Infusions to be performed under the close supervision of a healthcare professional and to observe the patient for least one hour after completion of the infusion. <input type="checkbox"/> P[{ ^A } ~• } Å~ æA T ÖU-æ	

By signing below, I certify that above therapy is medically necessary. Prescriber's Signature (SIGN BELOW)			
_____	_____	_____	_____
Dispense as Written	Date	Substitution Allowed	Date

The information contained in this facsimile may be confidential and is intended solely for the use of the named recipient(s). Access, copying or re-use of the facsimile or any information contained therein by any other person is not authorized. If you are not the intend recipient, please notify us immediately.