

Ocrevus
Referral Form

Y Medical Associates
Fax Referral To: 855-838-0623
Phone: 800-447-7558



Date: _____

PATIENT INFORMATION

Patient Name: _____
Address: _____
City, State, Zip: _____
Home Phone: _____
Cell Phone: _____
Date of Birth: _____
Gender: M F

PRESCRIBER INFORMATION

Prescriber Name: _____
Address: _____
City, State, Zip: _____
Phone: _____
Fax: _____
DEA#: _____ NPI#: _____
Contact Person: _____

INSURANCE INFORMATION (Please attach the front and back of insurance and prescription drug card)

Primary Insurance: _____ ID#: _____ Group: _____
Secondary Insurance: _____ ID#: _____ Group: _____
Prescription Card: _____ ID#: _____ BIN: _____ PCN: _____ Group: _____

DIAGNOSIS (ICD-10)

- G35 Relapsing forms of Multiple Sclerosis (Clinically isolated syndrome/relapsing-remitting disease/active secondary progressive disease)
- G35 Primary Progressive Multiple Sclerosis

PRE-SCREENING

- Hepatitis B Surface Antigen: _____
- Total Hepatitis B Core Antibody (Anti-HBc): _____
- Serum Immunoglobulins: _____

Vaccinations: Live-attenuated or live vaccines is not recommended during treatment and after discontinuation until B-cell repletion. Administer all necessary immunizations according to immunization guidelines at least 4 weeks prior to initiation for live or attenuated vaccines and at least 2 weeks prior to initiation for non-live vaccines.

Pre-screening: Required Hepatitis screening before first dose to include:

- Vaccination: _____ Height: _____ Weight: _____ Allergies: _____
- Hepatitis B Surface Antigen (HBsAg) and Total Hepatitis B Core Antibody (anti-HBc) * Ocrevus® is contraindicated in patients with active HBV. Patients who are negative for surface antigen HBsAg (-) and positive for HB core antibody HBcAB (+) or positive for surface antigen HBsAg (+), should consult liver disease experts before starting and during treatment.
- Quantitative Serum Immunoglobulin Screening (IgG, IgA, IgM)

(live or live-attenuated 4 weeks before, non-live 2 weeks before initiation of therapy)

Labs (During Therapy): _____

PRESCRIPTION ORDERS

Premeds

Premedication to be given 30 minutes prior to infusion:
 Acetaminophen PO: 325mg 500mg 650mg
 Diphenhydramine: 25mg IVP 50mg IVP 25mg PO 50mg PO
OR Alternate oral antihistamine: Cetirizine 10mg Loratadine 10mg
 Fexofenadine 60mgs Fexofenadine 180mgs

IV Access Flush Order: NaCl 0.9% 5-10ml IV before and after infusion

- Methylprednisolone 125mg IVP 40mg IVP OR _____mg PO
- Others/Miscellaneous: _____

Anaphylaxis Orders and Medications

Diphenhydramine Administer 25 mg slow IV/IM may repeat x1 **Dispense:** 1 x 50 mg vial

Epinephrine Autoinjector Administer 0.15mg (1:2000) IM (< 30 Kg)
 Administer 0.3mg (1:1000) IM (≥ 30 Kg)

Dispense: 1 package (2 pens)

Sodium Chloride 0.9% *Use to maintain IV line, prevent or treat hypotension in case of anaphylaxis*

Dispense: QS

Medication

Ocrevus (Ocrelizumab) IV as directed to infuse per protocol via pump with 0.22 µ [1:100] filter, following each infusion with a one hour post observation period.

Induction/Initial dosing: Induction/Initial dosing: 300mg Ocrevus IV in 250ml Sodium Chloride 0.9% to be infused at Week 0 over 2.5 hours or longer and 2 weeks later over 2.5 hours or longer. No Refills. ****To be infused in MD office or an Infusion suite.**

Maintenance dosing: 600mg Ocrevus IV in 500ml Sodium Chloride 0.9% to be infused every 6 months. 2 hrs or longer for eligible patients who have not experienced a serious infusion reaction with any previous Ocrevus Infusion 3.5-4 hrs or longer. Refills: X1 year ****Infusions to be performed under the close supervision of a healthcare professional and to observe the patient for least one hour after completion of the infusion.**

P[{ ^ Á } ~ • q } Á Ú x Á T Ö Á ~ Á

By signing below, I certify that above therapy is medically necessary. **Prescriber's Signature (SIGN BELOW)**

Dispense as Written

Date

Substitution Allowed

Date