## Y Medical Associates Crohn's and Ulcerative Colitis **YMedical Referral Form** Fax: 855-838-0623 Phone: 800-447-7558 ASSOCIATES Date: **Prescriber Information Patient Information** Prescriber Name: Please complete the following or send patient demographic sheet Address: Patient Name: City, State, Zip: Address: Phone: Fax: City, State, Zip: NPI#: DEA: Home Phone: Contact Person: Cell Phone: Gender: □ M DOB: $\Box F$ **Insurance Information** Primary Insurance:\_ ID#: Group:\_\_\_ Secondary Insurance: ID#: Group:\_\_\_ ID#: BIN#: PCN#: Prescription Card: Group: Medical Information (Section must be completed to process prescription) (Attach separate sheet if needed) Prior Authorization Insurance Number: Diagnosis - Please include diagnosis name with ICD-10 code **Therapy Details**: ☐ New ☐ Reauthorization ☐ Restart $\square$ K50.00 Crohn's disease of small intestines without complications Weight\_\_\_\_\_kg/lbs Height\_\_\_\_cm/in ☐ K50.8 Crohn's disease of both intestines without complications Allergies\_\_\_\_\_ ☐ K50.10 Crohn's disease of large intestines without complications Lab Data ☐ K50.00 Crohn's disease, unspecified, without complications Prior Therapies \_\_\_\_ ☐ K20.0 Eosinophilic Esophagitis Concomitant Medications \_\_\_\_\_ ☐ Other diagnosis: ICD-10 code Additional Comments Injection Training Required? Description \_\_\_Date of Description \_\_\_ ☐ Yes ☐ No Has a TB test been performed? □Yes □ No Does the Patient have an active infection? ☐ Yes □ No Start Date Review Date **Prescription Information** Medication Dose Strength Directions Qty Refills ☐ 200mg/mL Vial Kit ☐ 200 mg/mL Starter Ki ☐ Cimzia ☐ Loading Dose: Inject 400mg SC at Weeks 0, 2, and 4 ☐ 200mg/mL prefilled Syringe ☐ **Maintenance Dose:** Inject 200mg SC every 2 weeks ☐ PFS with needle shield 300 mg/2 mL □ Dupixent ☐ Inject 300 mg Sub Q every week ☐ Prefilled Pen 300 mg/2 mL ☐ Loading Dose: Inject 300mg IV over 30 minutes at ☐ Entyvio ☐ 300mg vial Weeks 0, 2, and 6. ☐ Maintenance Dose: Infuse 300mg IV over 30 minutes every 8 weeks Adult: Starter Kits: ☐ Humira $\square$ 80mg/0.8mL Starter Pack Pre-Filled Pen (Citrate Free) ☐ **Loading Dose:** Inject 160mg SC on Day 1, then 80mg ☐ Adalimumab on Day 15 (two weeks later) Maintenance: (biosimilar) ☐ 40mg/0.4mLPre-FilledPen(Citrate Free) ☐ **Maintenance Dose:** Inject 40mg SC every other week (starting Day 29) ☐ 40mg/0.4mLPre-FilledSyringe (Citrate Free) Pediatric (>6 years and adolescents) 17kg to < 40kg ☐ Other: ☐ **Loading Dose:** Inject 80mg SC on Day 1, 40mg on Day 15 (two weeks later) ☐ **MaintenanceDose:**Inject20mgSCeveryotherweek (starting Day 29) Pediatric (>6 years and adolescents) > 40kg

The information contained in this facsimile may be confidential and is intended solely for the use of the named recipient(s). Access, copying or re-use of the facsimile or any information contained therein by any other person is not authorized. If you are not the intend recipient, please notify us immediately by calling or faxing back to the originator.

DAW (Dispense as Written)

Prescriber Signature:\_

(starting Day 29)

☐ **Loading Dose:** Inject 160mg SC on Day 1, 80mg on

☐ **Maintenance Dose:** Inject 40mg SC every other week

Day 15 (two weeks later)

## Crohn's and Ulcerative Colitis Y Medical Associates **YMedical** Referral Form Fax: 855-838-0623 Phone: 800-447-7558 ASSOCIATES Date: **Patient Information Prescriber Information** Prescriber Name: Please complete the following or send patient demographic sheet Address: Patient Name: \_\_\_\_ City, State, Zip: Address: Phone: City, State, Zip: Fax: \_\_\_\_\_ Home Phone: NPI#: DEA: Cell Phone: Contact Person: Gender: □ M □ F DOB: Insurance Information ID#:\_\_\_\_\_ Primary Insurance:\_\_\_\_ \_Group:\_\_\_\_ Secondary Insurance: ID#: Group: ID#:\_\_\_\_ BIN#:\_\_\_\_\_PCN#:\_\_\_ Prescription Card:\_\_\_ Group: \_\_\_ Medical Information (Section must be completed to process prescription) (Attach separate sheet if needed) Prior Authorization Insurance Number: \_\_ **Diagnosis -** Please include diagnosis name with ICD-10 code **Therapy Details**: ☐ New ☐ Reauthorization ☐ Restart ☐ K50.00 Crohn's disease of small intestines without complications Weight\_\_\_\_kg/lbs Height\_\_\_\_cm/in Allergies\_\_\_\_\_ ☐ K50.8 Crohn's disease of both intestines without complications ☐ K50.10 Crohn's disease of large intestines without complications $\square$ K50.00 Crohn's disease, unspecified, without complications Prior Therapies \_\_\_\_ ☐ Other diagnosis: ICD-10 code Concomitant Medications Description Date of Description Additional Comments Injection Training Required? $\Box$ Yes $\Box$ No Has a TB test been performed? □ No □Yes Does the Patient have an active infection? ☐ Yes □ No Start Date Review Date **Prescription Information Dose Strength** Medication Directions Qty | Refills ☐ Avsola ☐ 100mg Vial ☐ **Loading Dose:** Infuse 5mg/kg at Weeks 0, 2, and 6 ☐ Inflectra ☐ Maintenance Dose: Infuse 5mg/kg every 8 weeks ☐ Remicade ☐ Renflexis ☐ Induction Therapy – 45 mg tablet ☐ **Induction Therapy:** 45 mg PO daily x 8 weeks. ☐ Rinvoq ☐ **Maintenance Therapy** – 15 mg or Maintenance Therapy: 30 mg tablets ☐ 15 mg PO daily ☐ 30 mg PO daily ☐ 100mg/mL SmartJect Auto Injector ☐ Simponi ☐ **Loading Dose:** Inject 200 mg SQat Week 0 then □100mg/mL Prefilled Syringe 100mg at Week 2 ☐ **Maintenance Dose:** Inject 100mg SQ every 4 weeks ☐ Stelara ☐ 130mg/26mL solution single dose vial $\square$ Loading Dose: Infuse: $\square$ 250mg $\square$ 390mg $\square$ 520mg ☐ 90mg/mL Prefilled Syringe as initial IV dose as directed by prescriber Date of Initial Infusion: \_\_\_\_ ☐ Maintenance Dose: Inject 90mg SC every 8 weeks (begin dosing 8 weeks after the IV induction dose) ☐ Initiation Therapy – 600 mg/10 mL single ☐ Initiation Therapy — ☐ Inject 600 mg IV over at least 1 ☐ Skyrizi nour at Weeks 0, 4, 8. 1vial/week. ☐ **Ongoing Therapy** – Week 12 – Inject 360 mg SC and every 8 weeks thereafter. 1 device with prefilled cartridge. ☐ Ongoing Therapy – 360 mg/2.4 mL prefilled cartridge with On-Body Injector ☐ Loading Dose: ☐ 5mg tablet ☐ 10mg twice daily for 8 weeks ☐ Xeljanz ☐ 10mg tablet ☐ XR: 22mg once for 8 weeks ☐ 11mgXRtablet ☐ **Maintenance Dose:** ☐ 5mg twice daily ☐ XR: 11mg ☐ 22mgXRtablet once daily ☐ 10mg twice daily ☐ XR: 22mg once daily \_\_\_\_\_DAW (Dispense as Written) Prescriber Signature: □Y □N Date: \_\_