IG and General Immune Disorders Enrollment Form	Fax: 8	lical Associates 855-838-0623 e: 800-447-7558	YMedical ASSOCIATES
Patient Name: Address: City, State, Zip: Home Phone: Cell Phone: Date of Birth:		Prescriber Name:	
INSURANCE INFORMATION (Primary Insurance: Secondary Insurance: Prescription Card:	Please attach the fro	ID#:	Group
DIAGNOSIS (ICD-10) Neurological G61.81 Chronic Inflammatory Demyelinating Polyneuropathy (CIDP) G61.82 Multifocal Motor Neuropathy (MMN) G61.0 Guillain-Barre G35 Multiple Sclerosis G70.01 Myasthenia Gravis w/Exacerbation Other:		Immunological Primary Immune Deficiency – Please specify ICD-10 Code: D80.9 Deficiency of Humoral Immunity D83.9 Common Variable Immunodeficiency D89.9 Immune Mechanism Disorder D81.9 Immune Deficiency NOS D69.3 Idiopathic Thrombocytopenia D80.1 Hypogammaglobulinemia Other:	
CLINICAL INFORMATION (Please a PatientWeight:Kg/Lbs Height: Has patient previously received IG Yes Medication	Inches/CM Allergies	nation, lab results, and other n s: ess:	
Intravenous IVIg Pharmacy Recommendation	(Pharmacy to round to Infuse total dose OVER week(s) for:		Infuse total dose of Immunoglobulin intravenously based on manufacturer recommend infusion rate as tolerated. Infuse via:
		s \Box 6 months \Box 12 months	Infusion Pump Excludes Medicare D
Medication	Other:	s 🗆 6 months 🗆 12 months	Excludes Medicare D
Medication Subcutaneous	Other: Dose grams OF (Pharmacy to round to Infuse total dose OVER week(s) for: 1 month	Rgram(s) perkg nearest vial size)	
Subcutaneous	Other: Dosegrams OF (Pharmacy to round to Infuse total dose OVERweek(s) for: 1 month 3 month Other nine (recommended) n: dose mtocol) and PRN	Rgram(s) perkg nearest vial size) day(s); Every s □ 6 months □ 12 months Diphenhydramine Administer 2 Dispense: 1 x 50 mg vial Epinephrine □ Administer 0.3m □ Administer 0.15r Dispense: 1 package	Excludes Medicare D Directions Infuse total dose of Immunoglobulin subcutaneously in one or more infusion sites via infusion pump based on manufacturer recommend infusion rate as tolerated. Other: Orders and Medications 5 mg slow IV/IM may repeat x1
Subcutaneous SCIg Pharmacy Recommendation Labs baseline and then every 6 months: BUN/Creating Premedication to be given 30 minutes prior to infusion Diphenhydramine IV or PO 25 mg or 50 mg Please circle route and dose Acetaminophen 325mg or 650 mg Please circle dose Ketorolac 30mg Slow IV-Push – 30mg/ml vial #1 per Other: IVAccess Flush Order: (Infusion supplies per pharmacy provide) NaCI 0.9% 5-10ml IV before and after infusion Heparin 10 units/ml 3-5ml IV after infusion for PICC/Midline at Heparin 100 units/ml 3-5ml IV after infusion for Port and PRN	Other: Dose grams OF (Pharmacy to round to Infuse total dose OVERweek(s) for: 1 month 3 month Other nine (recommended) n: dose btocol) and PRN tion ally necessary. Prescrib	gram(s) perkg nearest vial size) day(s); Every s 6 months 12 months Diphenhydramine Administer 2 Dispense: 1 x 50 mg vial Epinephrine Administer 0.3m Dispense: 1 package Sodium Chloride 0.9% Use to n case of anaphylaxis Dispense: QS	Excludes Medicare D Directions Infuse total dose of Immunoglobulin subcutaneously in one or more infusion sites via infusion pump based on manufacturer recommend infusion rate as tolerated. Other: Other: Orders and Medications 5 mg slow IV/IM may repeat x1 g (1:1000) Sub-Q (≥ 30 Kg) ng (1:2000) Sub-Q (< 30 Kg) maintain IV line, prevent or treat hypotension in

The information contained in this facsimile may be confidential and is intended solely for the use of the named recipient(s). Access, copying or re-use of the facsimile or any information contained therein by any other person is not authorized. If you are not the intend recipient, please notify us immediately.