Rheumatology	<b>Referral Form</b>
--------------	----------------------

Date: \_\_\_\_\_

## Y Medical Associates Fax:855-838-0623 Phone: 800-447-7558



	Patient Information	Prescriber Information	0.0	
Datiant Nama				
		Prescriber Name:		
	,.			
		Fax:NPI #:		
Gender: □M		Contact Person:		
		urance Information		
	urance: ID#:			
Prescription Car				
		e fax all pertinent clinical and lab information)		
	, , , , ,	ic Arthritis) □ L40.59 (Psoriatic Arthritis) □ L40.54 (Psoriat		
	osing Spondylitis) 🛛 M32.9 (Systemic Lupu	is Erythematosis)		
Diagnosis Date:		_		
		ssessment (Fill in below or attach lab work)		
		umber of Tender Joints: CRP:[		
		Current Height:Date:ESR:		
		Therapy Change   Stop Date:		
	• •	$\Box Weeks Completed: \Box 0 \Box 2$	□ 4	□ 6
	4. (Disease and its server af a servite)			
		- 14 - 1		
-	Score & Date (Please provide a copy of resu			
Medication	Dose Strength	Directions	Qty	Refills
Actemra	Prefilled Syringe 162mg/0.9mL     Auto Injector 162mg/0.9mL	□ <100kg Inject 162mg/0.9mL SC every 2 weeks □ >100kg Inject 162mg/0.9mL SC every week		
Benlysta	🗆 10mg/kg	□ IV Starter Dose: Infuse 10mg/kg every 2 weeks for		
		3 doses		
	200mg Autoinjector	□ IV Maintenance: Inject 10mg/kg every 4 weeks □ Inject 200mg SC once weekly (if switching from IV		
		administer first SC dose 1-4 weeks after last IV dose)		
Cimzia	🗆 Starter Kit 🗆 Syringe 🗆 Vial	□ Starter Dose: Inject 400mg SC on week 0, 2, and 4		
		□ Maintenance Dose: Inject 200mg SC every 2 weeks		
		□ Maintenance Dose: Inject 400mg SC once a month		
Cosentyx	□ 150mg Sensoready Pen	□ Starter Dose: Inject 150mg SC on week 0, 1, 2, 3,	□ 5	
	□ 300mg Sensoready Pen	and 4  Maintenance: Inject 150mg SC every 4 weeks	□ 1	
		Starter Dose: Inject 300mg SC on week 0, 1, 2, 3,	□ 10	
		and 4	-	
		□ Maintenance: Inject 300mg SC every 4 weeks	□2	
Enbrel	□ 25mg Syringe □ 0.25mg Vial	□ Inject 50mg SC every week		
	□ 50mg Syringe □ 50mg SureClick Pen □ Mini 50mg/mL	□ Injectmg(0.8mg/kg xkg) SC every week		
Evenity	□ 105mg/1.17mL	Inject 2 syringes (105mg each) for total dose of 210mg SQ once monthly	□2	
Forteo	□ 600mcg/2.4mL PFS	Maintenance: Inject 20mcg SC once daily	□1	
□ Humira	□ 10mg Syringe	□ Inject 10mg SC every other week (10 to <15kg)		
☐ Adalimumab	□ 20mg Syringe	□ Inject 20mg SC every other week (15 to <30kg)		
(biosimilar)	□ 40mg/0.4mL Syringe □ 40mg/0.4mL Pen	□ Inject 40mg SC every other week (30kg) □ Inject 40mg SC once weekly		
			I	I
Drossribar Sime	aturo	DAW (Dispanse as Writter) Date:		
Prescriber Sign	alui 6.	DAW (Dispense as Written) Date:		

The information contained in this facsimile may be confidential and is intended solely for the use of the named recipient(s). Access, copying or re-use of the facsimile or any information contained therein by any other person is not authorized. If you are not the intend recipient, please notify us immediately.

Date:

## Y Medical Associates Fax:855-838-0623 Phone: 800-447-7558



						-	
	Patient Information				riber Informa		
				<sup>r</sup> Name:			
				7:			
	D:		-	e, Zip:			
					NDI #		
Gender:				erson:			
		Incuran	ce Informatio				
					Creury		
-	ance:						
	surance: II rd: II			PCN#:			
-		mation (Please fax			-		
	matoid Arthritis)		-				(rthritic)
	losing Spondylitis) $\Box$ M32	•	, , ,	,	•		,
Diagnosis Date:				<u> </u>			
		and Clinical Asses	sment (Fill in b	elow or attach	lab work)		
Joints Affected	Bidgirosis					Date:	
	llen Joints:Cu						
	/ Induction   Stop Date:						
	ntinuation   Stop Date:						□ 6
TB Results & Da	ate (Please provide copy of	result):					
Bone Density	/ Score & Date (Please pro	vide a copy of results): _					
Medication	Dose Stren	gth		Directions		Qty	Refills
Kevzara	□ 150mg/1.14mL PFS	□ 200mg/1.14mL Pen	☐ Inject 200mg \$ ☐ Other:	SC once every 2	weeks		
Krystexxa	🗆 8mg/mL		□ Infuse 8mg in once every 2	250mL of NS ove 2 weeks	er 120 minutes		
Olumiant	□ 2mg Tablet	□ 1mg Tablet	🗆 Take one tabl	et PO daily			
Orencia	□ 125mg Pen □ 125mg Pen Syringe	□ 250mg Vial		fusemg veeks thereafter nject 125mg SC or	at weeks 0, 2, 4 nce a week		
Otezla	□ Starter Pack	□ 30mg Tablet	□ Starter Pack: □ Maintenance	Use as directed <b>Dose:</b> Take one	tablet PO BID		
Prolia	□ 60mg PFS		□ Inject 1 syring	e SC every 6 mor	nths		
<ul> <li>□ Avsola</li> <li>□ Inflectra</li> <li>□ Remicade</li> <li>□ Renflexis</li> </ul>	☐ 100mg Vial			e: Infuse 5mg/kg a Dose: Infuse 5m		6	
Rinvoq	□ 15mg		□ Take 1 tablet	PO daily			
☐ Rituxan ☐ Truxima	□ 100mg Vial	□ 500mg Vial	□ Infuse 1000m	ig on day 1 and d	ay 15		
Prescriber Sign	ature:		DAW (Di	spense as Writte	en) Date: _		

The information contained in this facsimile may be confidential and is intended solely for the use of the named recipient(s). Access, copying or re-use of the facsimile or any information contained therein by any other person is not authorized. If you are not the intend recipient, please notify us immediately.

Rheumatology Referral Form

Date:

## Y Medical Associates Fax:855-838-0623 Phone: 800-447-7558



	atient Information	Prescriber Ir		
Patient Name:		Prescriber Name:		
Address:		Address:		
City, State, Zip:		City, State, Zip:		
Home Phone:		Phone:		
DOB:		DEA:N	NPI #:	
Gender: □M □F		Contact Person:		
		Insurance Information		
Prescription Card:	ID#:		•	
	Clinical Information (Ple	ease fax all pertinent clinical and lab inforn	nation)	
M06.9 (Rheumatoid A	Arthritis) 🗆 M08.0 (Juvenile Idio	pathic Arthritis) 🗆 L40.59 (Psoriatic Arthritis) 🗆 L40	.54 (Psoriatic Juvenile	e Arthritis)
□ M45.9 (Ankylosing Sp	oondylitis) 🛛 M32.9 (Systemic L	_upus Erythematosis)		,
Diagnosis Date:	, ()			
	Diagnosis and Clinica	al Assessment (Fill in below or attach lab w	ork)	
Joints Affected:		Number of Tender Joints: CRP:		
		Ourrent Height:Date:		
		Outern HolginiOuter		
		Weeks Completed: □0		□ 6
Allergies:				
	se provide copy of result):			
	Date (Please provide a copy of			
Medication	Dose Strength	Directions	Qty	Refills
Simponi/Simponi Aria	<u>Simponi:</u>	<u>Simponi:</u>		
	$\Box$ SmartJect 50mg/0.5mL	☐ Inject 50mg SC once per month		
	□ 50mg/0.5mL PFS	Simponi Aria:		
	□ 50mg/0.5mL PFS <u>Simponi Aria:</u>	Simponi Aria: □ Infuse mg(2mg/kg) IV over 30		
Stelara	□ 50mg/0.5mL PFS <u>Simponi Aria:</u> □ 50mg/4mL Vial	Simponi Aria: Infuse mg(2mg/kg) IV over 30 minutes at 0 and 4 weeks, then every 8 weeks		
Stelara	□ 50mg/0.5mL PFS <u>Simponi Aria:</u> □ 50mg/4mL Vial □ 45mg/0.5mL PFS	Simponi Aria: Infuse mg(2mg/kg) IV over 30 minutes at 0 and 4 weeks, then every 8 weeks Inject 45mg SC on Day 1 (<100kg)		
Stelara	□ 50mg/0.5mL PFS <u>Simponi Aria:</u> □ 50mg/4mL Vial	Simponi Aria: Infuse mg(2mg/kg) IV over 30 minutes at 0 and 4 weeks, then every 8 weeks		
Stelara	□ 50mg/0.5mL PFS <u>Simponi Aria:</u> □ 50mg/4mL Vial □ 45mg/0.5mL PFS	Simponi Aria: □ Infuse mg(2mg/kg) IV over 30 minutes at 0 and 4 weeks, then every 8 weeks □ Inject 45mg SC on Day 1 (<100kg) □ Inject 90mg SC on day 1 (>100kg)		
Stelara	□ 50mg/0.5mL PFS <u>Simponi Aria:</u> □ 50mg/4mL Vial □ 45mg/0.5mL PFS	Simponi Aria:       mg(2mg/kg) IV over 30         Infuse       mg(2mg/kg) IV over 30         minutes at 0 and 4 weeks, then every 8 weeks         Inject 45mg SC on Day 1 (<100kg)		
Stelara	□ 50mg/0.5mL PFS <u>Simponi Aria:</u> □ 50mg/4mL Vial □ 45mg/0.5mL PFS	Simponi Aria: Infuse mg(2mg/kg) IV over 30 minutes at 0 and 4 weeks, then every 8 weeks Inject 45mg SC on Day 1 (<100kg) Inject 90mg SC on day 1 (>100kg) Inject 45mg SC on Day 29 and every 12 weeks thereafter (<100kg) Inject 90mg SC on Day 29 and every 12 weeks thereafter (>100kg)		
Stelara	<ul> <li>□ 50mg/0.5mL PFS</li> <li><u>Simponi Aria:</u></li> <li>□ 50mg/4mL Vial</li> <li>□ 45mg/0.5mL PFS</li> <li>□ 90mg/mL PFS</li> <li>□ 150 mg/mL in each single-</li> </ul>	Simponi Aria:       □         □ Infuse       mg(2mg/kg) IV over 30         minutes at 0 and 4 weeks, then every 8 weeks         □ Inject 45mg SC on Day 1 (<100kg)		
	<ul> <li>□ 50mg/0.5mL PFS</li> <li><u>Simponi Aria:</u></li> <li>□ 50mg/4mL Vial</li> <li>□ 45mg/0.5mL PFS</li> <li>□ 90mg/mL PFS</li> <li>□ 150 mg/mL in each single-dose prefilled pen</li> </ul>	Simponi Aria:         □ Infuse       mg(2mg/kg) IV over 30 minutes at 0 and 4 weeks, then every 8 weeks         □ Inject 45mg SC on Day 1 (<100kg)		
	<ul> <li>□ 50mg/0.5mL PFS</li> <li><u>Simponi Aria:</u></li> <li>□ 50mg/4mL Vial</li> <li>□ 45mg/0.5mL PFS</li> <li>□ 90mg/mL PFS</li> <li>□ 150 mg/mL in each single-dose prefilled pen</li> <li>□ 90 mg/mL in each single-</li> </ul>	Simponi Aria:       □         □ Infuse       mg(2mg/kg) IV over 30         minutes at 0 and 4 weeks, then every 8 weeks         □ Inject 45mg SC on Day 1 (<100kg)		
	<ul> <li>50mg/0.5mL PFS</li> <li><u>Simponi Aria:</u> <ul> <li>50mg/4mL Vial</li> <li>45mg/0.5mL PFS</li> <li>90mg/mL PFS</li> </ul> </li> <li>150 mg/mL in each single-dose prefilled pen</li> <li>90 mg/mL in each single-dose prefilled syringe</li> </ul>	Simponi Aria:         □ Infuse       mg(2mg/kg) IV over 30 minutes at 0 and 4 weeks, then every 8 weeks         □ Inject 45mg SC on Day 1 (<100kg)		
	<ul> <li>□ 50mg/0.5mL PFS</li> <li><u>Simponi Aria:</u></li> <li>□ 50mg/4mL Vial</li> <li>□ 45mg/0.5mL PFS</li> <li>□ 90mg/mL PFS</li> <li>□ 150 mg/mL in each single-dose prefilled pen</li> <li>□ 90 mg/mL in each single-</li> </ul>	Simponi Aria:         □ Infuse       mg(2mg/kg) IV over 30 minutes at 0 and 4 weeks, then every 8 weeks         □ Inject 45mg SC on Day 1 (<100kg)		
	<ul> <li>50mg/0.5mL PFS</li> <li><u>Simponi Aria:</u> <ul> <li>50mg/4mL Vial</li> <li>45mg/0.5mL PFS</li> <li>90mg/mL PFS</li> </ul> </li> <li>150 mg/mL in each single-dose prefilled pen</li> <li>90 mg/mL in each single-dose prefilled syringe</li> <li>150 mg/mL in each single-dose prefilled syringe</li> <li>150 mg/mL in each single-dose prefilled syringe</li> </ul>	Simponi Aria: Infuse mg(2mg/kg) IV over 30 minutes at 0 and 4 weeks, then every 8 weeks Inject 45mg SC on Day 1 (<100kg) Inject 90mg SC on day 1 (>100kg) Inject 45mg SC on Day 29 and every 12 weeks thereafter (<100kg) Inject 90mg SC on Day 29 and every 12 weeks thereafter (>100kg) Initial Dose: Inject 150mg SC weeks 0, and 4 Maintenance Dose: Inject 150mg SC every 12 weeks		
	<ul> <li>50mg/0.5mL PFS</li> <li><u>Simponi Aria:</u> <ul> <li>50mg/4mL Vial</li> <li>45mg/0.5mL PFS</li> <li>90mg/mL PFS</li> </ul> </li> <li>150 mg/mL in each single-dose prefilled pen</li> <li>90 mg/mL in each single-dose prefilled syringe</li> <li>150 mg/mL in each single-dose prefilled syringe</li> <li>150 mg/mL in each single-dose prefilled syringe</li> </ul>	Simponi Aria: Infusemg(2mg/kg) IV over 30 minutes at 0 and 4 weeks, then every 8 weeks Inject 45mg SC on Day 1 (<100kg) Inject 90mg SC on day 1 (>100kg) Inject 45mg SC on Day 29 and every 12 weeks thereafter (<100kg) Inject 90mg SC on Day 29 and every 12 weeks thereafter (>100kg) Initial Dose: Inject 150mg SC weeks 0, and 4 Maintenance Dose: Inject 150mg SC every 12 weeks Starter Dose: Inject 160mg SC on Day 1		
Skyrizi	<ul> <li>50mg/0.5mL PFS</li> <li><u>Simponi Aria:</u></li> <li>50mg/4mL Vial</li> <li>45mg/0.5mL PFS</li> <li>90mg/mL PFS</li> <li>150 mg/mL in each single-dose prefilled pen</li> <li>90 mg/mL in each single-dose prefilled syringe</li> <li>150 mg/mL in each single-dose prefilled syringe</li> <li>150 mg/mL AutoInjector</li> </ul>	Simponi Aria: Infuse mg(2mg/kg) IV over 30 minutes at 0 and 4 weeks, then every 8 weeks Inject 45mg SC on Day 1 (<100kg) Inject 90mg SC on day 1 (>100kg) Inject 45mg SC on Day 29 and every 12 weeks thereafter (<100kg) Inject 90mg SC on Day 29 and every 12 weeks thereafter (>100kg) Initial Dose: Inject 150mg SC weeks 0, and 4 Maintenance Dose: Inject 150mg SC every 12 weeks		
Skyrizi Taltz	<ul> <li>50mg/0.5mL PFS</li> <li><u>Simponi Aria:</u></li> <li>50mg/4mL Vial</li> <li>45mg/0.5mL PFS</li> <li>90mg/mL PFS</li> <li>90mg/mL PFS</li> <li>150 mg/mL in each single- dose prefilled pen</li> <li>90 mg/mL in each single- dose prefilled syringe</li> <li>150 mg/mL in each single- dose prefilled syringe</li> <li>150 mg/mL AutoInjector</li> <li>100mg PFS</li> </ul>	Simponi Aria:       mg(2mg/kg) IV over 30         Infuse       mg(2mg/kg) IV over 30         minutes at 0 and 4 weeks, then every 8 weeks         Inject 45mg SC on Day 1 (<100kg)		
Skyrizi	<ul> <li>□ 50mg/0.5mL PFS</li> <li><u>Simponi Aria:</u></li> <li>□ 50mg/4mL Vial</li> <li>□ 45mg/0.5mL PFS</li> <li>□ 90mg/mL PFS</li> <li>□ 150 mg/mL in each single-dose prefilled pen</li> <li>□ 90 mg/mL in each single-dose prefilled syringe</li> <li>□ 150 mg/mL in each single-dose prefilled syringe</li> <li>□ 150 mg/mL AutoInjector</li> <li>□ 100mg PFS</li> <li>□ 100mg One-Press</li> </ul>	Simponi Aria: Infusemg(2mg/kg) IV over 30 minutes at 0 and 4 weeks, then every 8 weeks Inject 45mg SC on Day 1 (<100kg) Inject 90mg SC on day 1 (>100kg) Inject 45mg SC on Day 29 and every 12 weeks thereafter (<100kg) Inject 90mg SC on Day 29 and every 12 weeks thereafter (>100kg) Initial Dose: Inject 150mg SC weeks 0, and 4 Maintenance Dose: Inject 150mg SC every 12 weeks Starter Dose: Inject 160mg SC on Day 1 Maintenance: Inject 80mg SC every 4 weeks Inject SubQ 100 mg at weeks 0, 4, and then		
Skyrizi Taltz Tremfya	<ul> <li>50mg/0.5mL PFS</li> <li><u>Simponi Aria:</u></li> <li>50mg/4mL Vial</li> <li>45mg/0.5mL PFS</li> <li>90mg/mL PFS</li> <li>90mg/mL PFS</li> <li>150 mg/mL in each single- dose prefilled pen</li> <li>90 mg/mL in each single- dose prefilled syringe</li> <li>150 mg/mL in each single- dose prefilled syringe</li> <li>80mg/mL AutoInjector</li> <li>100mg PFS</li> <li>100mg One-Press autoinjector</li> </ul>	Simponi Aria:       mg(2mg/kg) IV over 30         Infuse       mg(2mg/kg) IV over 30         minutes at 0 and 4 weeks, then every 8 weeks         Inject 45mg SC on Day 1 (<100kg)		
Skyrizi Taltz	<ul> <li>□ 50mg/0.5mL PFS</li> <li><u>Simponi Aria:</u></li> <li>□ 50mg/4mL Vial</li> <li>□ 45mg/0.5mL PFS</li> <li>□ 90mg/mL PFS</li> <li>□ 150 mg/mL in each single-dose prefilled pen</li> <li>□ 90 mg/mL in each single-dose prefilled syringe</li> <li>□ 150 mg/mL in each single-dose prefilled syringe</li> <li>□ 150 mg/mL AutoInjector</li> <li>□ 100mg PFS</li> <li>□ 100mg One-Press</li> </ul>	Simponi Aria:       mg(2mg/kg) IV over 30         Infuse       mg(2mg/kg) IV over 30         minutes at 0 and 4 weeks, then every 8 weeks         Inject 45mg SC on Day 1 (<100kg)	□ 1-Prefilled Pen	
Skyrizi Taltz Tremfya	<ul> <li>50mg/0.5mL PFS</li> <li><u>Simponi Aria:</u></li> <li>50mg/4mL Vial</li> <li>45mg/0.5mL PFS</li> <li>90mg/mL PFS</li> <li>90mg/mL PFS</li> <li>150 mg/mL in each single- dose prefilled pen</li> <li>90 mg/mL in each single- dose prefilled syringe</li> <li>150 mg/mL in each single- dose prefilled syringe</li> <li>80mg/mL AutoInjector</li> <li>100mg PFS</li> <li>100mg One-Press autoinjector</li> </ul>	Simponi Aria:       mg(2mg/kg) IV over 30         Infuse       mg(2mg/kg) IV over 30         minutes at 0 and 4 weeks, then every 8 weeks         Inject 45mg SC on Day 1 (<100kg)	□ 1-Prefilled Pen	
Skyrizi Taltz Tremfya Tymlos	<ul> <li>50mg/0.5mL PFS</li> <li><u>Simponi Aria:</u></li> <li>50mg/4mL Vial</li> <li>45mg/0.5mL PFS</li> <li>90mg/mL PFS</li> <li>90mg/mL PFS</li> <li>150 mg/mL in each single-dose prefilled pen</li> <li>90 mg/mL in each single-dose prefilled syringe</li> <li>150 mg/mL in each single-dose prefilled syringe</li> <li>150 mg/mL in each single-dose prefilled syringe</li> <li>100mg/mL AutoInjector</li> <li>100mg PFS</li> <li>100mg One-Press autoinjector</li> <li>80mcg/0.04mL</li> </ul>	Simponi Aria:       mg(2mg/kg) IV over 30         Infuse       mg(2mg/kg) IV over 30         minutes at 0 and 4 weeks, then every 8 weeks         Inject 45mg SC on Day 1 (<100kg)	□ 1-Prefilled Pen	
Skyrizi Taltz Tremfya	<ul> <li>□ 50mg/0.5mL PFS</li> <li><u>Simponi Aria:</u></li> <li>□ 50mg/4mL Vial</li> <li>□ 45mg/0.5mL PFS</li> <li>□ 90mg/mL PFS</li> <li>□ 90mg/mL in each single-dose prefilled pen</li> <li>□ 90 mg/mL in each single-dose prefilled syringe</li> <li>□ 150 mg/mL in each single-dose prefilled syringe</li> <li>□ 150 mg/mL in each single-dose prefilled syringe</li> <li>□ 150 mg/mL AutoInjector</li> <li>□ 100mg PFS</li> <li>□ 100mg One-Press autoinjector</li> <li>□ 80mcg/0.04mL</li> <li><u>Xeljanz:</u></li> </ul>	Simponi Aria:       mg(2mg/kg) IV over 30         Infuse       mg(2mg/kg) IV over 30         minutes at 0 and 4 weeks, then every 8 weeks         Inject 45mg SC on Day 1 (<100kg)		
Skyrizi Taltz Tremfya Tymlos	<ul> <li>□ 50mg/0.5mL PFS</li> <li><u>Simponi Aria:</u></li> <li>□ 50mg/4mL Vial</li> <li>□ 45mg/0.5mL PFS</li> <li>□ 90mg/mL PFS</li> <li>□ 90mg/mL in each single-dose prefilled pen</li> <li>□ 90 mg/mL in each single-dose prefilled syringe</li> <li>□ 150 mg/mL in each single-dose prefilled syringe</li> <li>□ 150 mg/mL in each single-dose prefilled syringe</li> <li>□ 150 mg/mL AutoInjector</li> <li>□ 100mg PFS</li> <li>□ 100mg One-Press autoinjector</li> <li>□ 80mcg/0.04mL</li> </ul>	Simponi Aria:       mg(2mg/kg) IV over 30         Infuse       mg(2mg/kg) IV over 30         minutes at 0 and 4 weeks, then every 8 weeks         Inject 45mg SC on Day 1 (<100kg)	□ 1-Prefilled Pen	
Skyrizi Taltz Tremfya Tymlos	<ul> <li>□ 50mg/0.5mL PFS</li> <li><u>Simponi Aria:</u></li> <li>□ 50mg/4mL Vial</li> <li>□ 45mg/0.5mL PFS</li> <li>□ 90mg/mL PFS</li> <li>□ 90mg/mL in each single-dose prefilled pen</li> <li>□ 90 mg/mL in each single-dose prefilled syringe</li> <li>□ 150 mg/mL in each single-dose prefilled syringe</li> <li>□ 150 mg/mL in each single-dose prefilled syringe</li> <li>□ 150 mg/mL AutoInjector</li> <li>□ 100mg PFS</li> <li>□ 100mg One-Press autoinjector</li> <li>□ 80mcg/0.04mL</li> <li><u>Xeljanz:</u></li> <li>□ 5mg Tablet <u>Xeljanz XR:</u></li> </ul>	Simponi Aria:       mg(2mg/kg) IV over 30         Infuse       mg(2mg/kg) IV over 30         minutes at 0 and 4 weeks, then every 8 weeks         Inject 45mg SC on Day 1 (<100kg)	□ 60	
Skyrizi Taltz Tremfya Tymlos	<ul> <li>□ 50mg/0.5mL PFS</li> <li><u>Simponi Aria:</u></li> <li>□ 50mg/4mL Vial</li> <li>□ 45mg/0.5mL PFS</li> <li>□ 90mg/mL PFS</li> <li>□ 90mg/mL in each single-dose prefilled pen</li> <li>□ 90 mg/mL in each single-dose prefilled syringe</li> <li>□ 150 mg/mL in each single-dose prefilled syringe</li> <li>□ 150 mg/mL in each single-dose prefilled syringe</li> <li>□ 150 mg/mL AutoInjector</li> <li>□ 100mg PFS</li> <li>□ 100mg One-Press autoinjector</li> <li>□ 80mcg/0.04mL</li> </ul>	Simponi Aria:       mg(2mg/kg) IV over 30         Infuse       mg(2mg/kg) IV over 30         minutes at 0 and 4 weeks, then every 8 weeks         Inject 45mg SC on Day 1 (<100kg)		

Prescriber Signature: \_\_\_\_\_ DAW (Dispense as Written) Date: \_\_\_\_\_

The information contained in this facsimile may be confidential and is intended solely for the use of the named recipient(s). Access, copying or re-use of the facsimile or any information contained therein by any other person is not authorized. If you are not the intend recipient, please notify us immediately.