IG and General Immune Disorders EnrolIment Form	Fax Referra	cal Associates al To: 855-838-0623 e: 800-447-7558	
PATIENT INFORMATION Patient Name: Address: City, State, Zip: Home Phone: Cell Phone: Date of Birth:		Prescriber Name:	RIBER INFORMATION
INSURANCE INFORMATION (A Primary Insurance: Secondary Insurance: Prescription Card:		ID#: ID#: BIN:PCN:PCN:	Group:
DIAGNOSIS (ICD-10) Neurological G61.81 Chronic Inflammatory Demyelinating Polyneurop G61.82 Multifocal Motor Neuropathy (MMN) G61.0 Guillain-Barre G35 Multiple Sclerosis G70.01 Myasthenia Gravis w/Exacerbation Other: CLINICAL INFORMATION (Please attempt of the second	aon Syndrome tach all clinical information ches/CM Allergies	 D80.9 Deficiency of Humoral Ir D83.9 Common Variable Immu D89.9ImmuneMechanism Disc D69.3 Idiopathic Thrombocytope Other: 	se specify ICD-10 Code: nmunity unodeficiency order
Medication Intravenous Intravenous IVIg Pharmacy Recommendation Infuse at home	(pharmacy to round to n Infuse total dose OVER grams per da Everyweeks for □ 1 month □ 3 months	days(s) OR ay fordays	Directions Infuse total dose of Immunoglobulin intravenously based on manufacturer recommend infusion rate as tolerated. Infuse via: Gravity Infusion Pump Excludes Medicare D
 Infuse at physician office Medication 	Other: Dose		Directions
Subcutaneous	(Pharmacy to round to m Infuse total dose OVER week(s) for:	day(s); Every s □ 6 months □ 12 months	Infuse total dose of Immunoglobulin subcutaneously in one or more infusion sites via infusion pump based on manufacturer recommend infusion rate as tolerated. Other:
Labs baseline and then every 6 months: BUN/Creatinine (recommended) Premedication to be given 30 minutes prior to infusion: Diphenhydramine IV or PO 25 mg or 50 mg Please circle route and dose Acetaminophen 325mg or 650 mg Please circle dose Ketorolac 30mg Slow IV-Push – 30mg/ml vial #1 per dose Other: IV Access Flush Order: (Infusion supplies per pharmacy protocol) NaCl 0.9% 5-10ml IV before and after infusion Heparin 10 units/ml 3-5ml IV after infusion for PICC/Midline and PRN Heparin 100 units/ml 3-5ml IV after infusion for POCC/Midline and PRN All infusion supplies necessary to administer the medication By signing below, I certify that above therapy is medically necessary. Prescribe		Anaphylaxis Orders and Medications Diphenhydramine Administer 25 mg slow IV/IM may repeat x1 Dispense: 1 x 50 mg vial Epinephrine □ Administer 0.3mg (1:1000) IM (≥ 30 Kg) □ Administer 0.15mg (1:2000) IM (< 30 Kg)	
 □ Ketorolac 30mg Slow IV-Push – 30mg/ml vial #1 per do □ Other:	col) d PRN RN on	Sodium Chloride 0.9% Use to m in case of anaphylaxis Dispense : QS	

The information contained in this facsimile may be confidential and is intended solely for the use of the named recipient(s). Access, copying or re-use of the facsimile or any information contained therein by any other person is not authorized. If you are not the intend recipient, please notify us immediately.